INDEX TO SCIENTIFIC NAMES

HERPETOLOGICA

Volume 36 December 1980

Acanthophis 281, 282, 285, 286
  antarcticus 281–287
Acrochordus 327
Adelotus brevis 264
Aegla 317
Aextoxicum punctatum 317
Afrixalus sp. 264
Agama agama 102
Agkistrodon contortrix 286
Alligator 289, 332
  mississippiensis 27, 289, 294, 364
  sclerops 363
Alsodes 316
Alytes 350
  obstetricans 277, 278, 350
Ambystoma 154, 156
  jeffersonianum 145
  macrodactylum 145
  maculatum 141, 145, 156
  mexicanum 46
  opacum 145
  tigrinum 47, 144–157
Amphibolurus muricatus 284
Amphisbaena cinerea 370
Andropogon hallii 128
  scoparius 128
Angiopteris evecta 3
Angolosaurus 312
Anolis baleatus 93
  barahonae 93–98
  barahonae albocellatus 94–98
  barahonae barahonae 94–98
  barahonae ***ininquinatus*** (subsp. nov.) 94–98
  barahonae ***mulitus*** (subsp. nov.) 95–98
  marmoratus 98
  oculatus 98
  recordi 93
  recordi subsolanus 98
Antechinus spp. 284
Aristida stricta 353
Arthroleptis sp. 264
Ascaphus 277–279, 346, 349–351
  truei 264, 275–279, 346–349
Asterophrys turpicula 264
Barygenes atra 264
  cheesmanae 264
Batagur baska 158
Batrachoseps attenuatus 14
Batrachyla 2, 3, 316
  antartandica 2
  leptopus 2, 317
  taeniata 317
Bitis arietans 170, 171
Blanus 370
Blechnum 317
Bolitoglossa 335, 336, 338, 339
  brevipes 336, 337
  franklini 335–344
  lincolni 336, 337
  nigroflavescens 336, 337
  resplendens 335–344
Bombina 133–137, 277, 278, 350, 365
  bombina 133, 135, 136, 264, 276, 278, 279
  orientalis 277–279
  variegata 133–135, 365
Bothrops 231
  caribbaea 231
  schlegeli 231–233
Breviceps 2, 263
  mossambica 258
  mossambicus 258, 264
  poweri 264
  sp. 264
  verrucosus 260
Bromus rubens 303
Bufo 136, 234, 316, 365
  americanus 234–236, 238–241
  boreas 277, 278
  bufo 37, 39–41
  bufo andrewsi 41
  bufo bankorensis 40
  bufo bufo 40
  bufo gargarizans 39–41
  cognatus 277, 278
  exsul 234
  kangi 37, 39, 40
  marinus 234, 264, 277, 278
  orientalis 39
  rangeri 277, 278
  rosei 264
  spinulosus 277, 278
  stejnegeri 37, 39–41
  stomaticus 39
  terrestris 234
  torrenticola 41
  viridus 39, 40
  woodhousei 234, 235, 238–242
Cacosternum sp. 264
Caiman 361
  crocodilus 361–363
Calamovilfa longifolia 128
Callisaurus draconoides 327
Calotes versicolor 116–118
Caretta caretta 18, 33, 34, 36, 72, 163
Caudiverbera 316
Centrolenella 318
  fleischmanni 318–320, 322, 324
  viridissima 324
Ceratobatrachus guentheri 264
Chamaesaura 312

INDEX TO SCIENTIFIC NAMES

HERPETOLOGICA

Volume 36 December 1980

Acanthophis 281, 282, 285, 286
  antarcticus 281–287
Acrochordus 327
Adelotus brevis 264
Aegla 317
Aextoxicum punctatum 317
Afrixalus sp. 264
Agama agama 102
Agkistrodon contortrix 286
Alligator 289, 332
  mississippiensis 27, 289, 294, 364
  sclerops 363
Alsodes 316
Alytes 350
  obstetricans 277, 278, 350
Ambystoma 154, 156
  jeffersonianum 145
  macrodactylum 145
  maculatum 141, 145, 156
  mexicanum 46
  opacum 145
  tigrinum 47, 144–157
Amphibolurus muricatus 284
Amphisbaena cinerea 370
Andropogon hallii 128
  scoparius 128
Angiopteris evecta 3
Angolosaurus 312
Anolis baleatus 93
  barahonae 93–98
  barahonae albocellatus 94–98
  barahonae barahonae 94–98
  barahonae ***ininquinatus*** (subsp. nov.) 94–98
  barahonae ***mulitus*** (subsp. nov.) 95–98
  marmoratus 98
  oculatus 98
  recordi 93
  recordi subsolanus 98
Antechinus spp. 284
Aristida stricta 353
Arthroleptis sp. 264
Ascaphus 277–279, 346, 349–351
  truei 264, 275–279, 346–349
Asterophrys turpicula 264
Barygenes atra 264
  cheesmanae 264
Batagur baska 158
Batrachoseps attenuatus 14
Batrachyla 2, 3, 316
  antartandica 2
  leptopus 2, 317
  taeniata 317
Bitis arietans 170, 171
Blanus 370
Blechnum 317
Bolitoglossa 335, 336, 338, 339
  brevipes 336, 337
  franklini 335–344
  lincolni 336, 337
  nigroflavescens 336, 337
  resplendens 335–344
Bombina 133–137, 277, 278, 350, 365
  bombina 133, 135, 136, 264, 276, 278, 279
  orientalis 277–279
  variegata 133–135, 365
Bothrops 231
  caribbaea 231
  schlegeli 231–233
Breviceps 2, 263
  mossambica 258
  mossambicus 258, 264
  poweri 264
  sp. 264
  verrucosus 260
Bromus rubens 303
Bufo 136, 234, 316, 365
  americanus 234–236, 238–241
  boreas 277, 278
  bufo 37, 39–41
  bufo andrewsi 41
  bufo bankorensis 40
  bufo bufo 40
  bufo gargarizans 39–41
  cognatus 277, 278
  exsul 234
  kangi 37, 39, 40
  marinus 234, 264, 277, 278
  orientalis 39
  rangeri 277, 278
  rosei 264
  spinulosus 277, 278
  stejnegeri 37, 39–41
  stomaticus 39
  terrestris 234
  torrenticola 41
  viridus 39, 40
  woodhousei 234, 235, 238–242
Cacosternum sp. 264
Caiman 361
  crocodilus 361–363
Calamovilfa longifolia 128
Callisaurus draconoides 327
Calotes versicolor 116–118
Caretta caretta 18, 33, 34, 36, 72, 163
Caudiverbera 316
Centrolenella 318
  fleischmanni 318–320, 322, 324
  viridissima 324
Ceratobatrachus guentheri 264
Chamaesaura 312

Chelonia depressa 269
mydas 17, 19, 72, 161–163
Chelydra 33, 35, 36
serpentina 33, 36, 158, 159, 360
Chiromantis xerampelina 264
Choerophryne rostellifer 264
Chrysemys picta 103, 158
Chrysobatrachus 3
cupreonitens 2
Clemmys guttata 158
Cnemidophorus sexlineatus 128
Colostethus 189, 196, 199
collaris 189, 195, 198, 199
inguinalis 189, 197–199
palmatus 189, 199
trinitatis 189–195, 197–199
Coluber 22–25
constrictor 20
Cophixalus 200
neglectus 264
pansus 264
riparius 258, 264
verrucosus 264
Copiula oxyrhina 264
Cordylosaurus 312
Cordylus 312, 315
cataphractus 311–315
cordylus 311–313
cordylus jonesii 311
cordylus rhodesianus 311
giganteus 311–315
polyzonus 311–313
vittifer 311–313
warreni 311–313
Coronella 92
austriaca 92
Crinia georgiana 264
Crocodilurus 314
Crocodylus acutus 364
Crotalus 326
atrox 327–332
bothrops 292
lepidus 120, 327–331
molossus 327, 328, 330, 331
ruber 120
scutulatus 327, 328, 330, 331
viridis 327, 328, 330, 331
Cryptoblepharus plagiocephalus 284
Cryptomeria japonica 305
Ctenotus robustus 284
taeniolatus 284
Cyclorana 257
australis 258, 264
longipes 264
Cynops pyrrhogaster 47
Cyrtodactylus 66–68
biordinis (sp. nov.) 66–69
loriae 66, 67
louisiadensis 66–69
novaeguineae 66, 67
pelagicus 66, 68
Dasypus novemcinctus 358
Dendrobates 178
auratus 70, 71
azureus 70, 71
femoralis 178, 179
granuliferus 1, 70, 71
histrionicus 70, 71
pumilio 69–71
Dermochelys coriacea 18
Dermophis 244, 246, 249
mexicanus 244–246, 249–255
Desmognathus 14, 113
fuscus 14, 109–114
ochrophaeus 109–114, 143
quadramaculatus 78, 85
Didelphis virginianus 358
Dieffenbachia 319, 320, 325
Diplostomulum 47
Dipsosaurus dorsalis 327
Discodeles sp. 264
Discoglossus 350
pictus 277, 278, 350
Drosophila melanogaster 295
Elachistocleis 258
sp. 258, 264
Elaphe 22–25, 230, 326
guttata emoryi 207
obsoleta 20, 120, 126
subocularis 327–331
Eleutherodactylus 175, 177, 179
achatinus 175
anomalus 175
areolatus 175, 179
biporcatus 175
brederi 175
bufoniformis 175
calcaratus 181
caprifer 175
caryophyllaceus 175, 185
chalceus 175, 179, 180, 185
coqui 277, 278
crucifer 181
cruentus 175, 182
diastema 175
dubitus 175
fitzingeri 175–178
florulentus 175
frater 184, 185
gaigeae 175–179, 187
gularis 175
lacrimosus 179
latidiscus 175
lehmanvalenciae 175
longirostris 175
moro 175
ockendeni 180, 184, 185
phoxocephalus 183
planirostris 232
quaquaversus 184, 185
raniformis 175
ranoides 175
ridens 175, 180, 181, 187
roseus 175, 180–182, 187

rugulosus 178
sp. A 175
sp. Z 175
subsigillatus 175, 182–184, 187
taeniatus 175, 180, 183–187
unistrigatus 176, 179
Emys orbicularis 33, 36, 136
Eretmochelys 265
imbricata 18, 163, 265, 269
Eumeces 202
multivirgatus 128
Eupsophus 316
migueli 317
roseus 317
vittatus 317
Eurycea 51, 53, 57–59, 113
aquatica 57, 58
bislineata 51–54, 57, 78, 109–114, 140
junaluska 51, 52
longicauda 51, 53, 59
lucifuga 51, 53, 59
multiplicata 51, 52, 59
neotenes 58
quadridigitata 51, 52, 138–140
tynerensis 51–54, 57–59
Farancia abacura 331
Gastrophryne 2, 234, 277
carolinensis 234, 235, 238–242, 277–279
Gastrotheca cavia 276, 278
riobambae 276, 278
Genyophryne thomsoni 264
Geochelone elephantopus 359
gigantea 302, 356, 359, 360
Geophis nasalis 255
Geotrypetes grandisonae 253
seraphini 253
Gerrhonotus 202
Gerrhosaurus 312, 314
flavigularis 311
major 311–313
validus 311–313
Gopherus 301, 353
agassizii 301–304, 360
polyphemus 60, 64, 301, 353, 356, 360
Gymnopis multiplicata 249
Gyrinophilus porphyriticus 78–86
Heleioporus eyrei 261, 264
Heleophryne purcelli 264
Hemidactylium scutatum 141–143
Hemisus marmoratus 264
Heterodon 136, 331
platyrhinos 169
Holbrookia 129–131
maculata 127–132
Hyla 257, 365
arborea savignyi 276, 278
cinerea 232, 276, 278
eximia 277, 278
femoralis 276, 278
gratiosa 276, 278
regilla 234
septentrionalis 234
Hylodes roseus 180
subsigillatus 182
taeniatus 184
Hylophorbus rufescens 264
Hylorina 316
Hymenophyllum 317
Hyperolius marmoratus 264
Hypopterigium arbuscula 317
Iguana 221
iguana 65, 221, 222, 224
Insuetophrynus 316
acarpicus 316–318
Kalmia latifolia 79
Kaloula 2, 277
conjuncta 264
pulchra 264, 276–279
Kassina sp. 264
Koeleria crisata 128
Lacerta agilis 60, 61, 64
Lampropeltis getulus 120, 126
mexicana 120
Lampropholis spp. 284
Lanchesis muta 42
Langloisia setosissima 302
Lechriodus 261
fletcheri 258, 264
melanopyga 264
Leiolopisma spp. 284
Leiopelma 346, 349–351
archeyi 349
hamiltoni 349
hochstetteri 264, 349, 350
Lepidochelys olivacea 18
Leptodactylus albilabris 234
pentadactylus 276, 278
sibilator 264
Leptopelis sp. 264
Limnodynastes dumerili 261, 264
ornatus 264
peroni 261, 264
spenceri 264
tasmaniensis 264
Lissemys punctata 102
Lithodytes 176
lineatus 177–179
Litoria 257
alboguttata 264
caerulea 264
chloris 264
dahlii 264
ewingi 264
fallax 264
freycinetti 264
infrafrenata 262, 264
lesueuri 264
nasuta 264
peroni 264
raniformis 264
rothi 264
rubella 264
wotjulumensis 264
Lodoicea maldivica 217

Mabuya 99, 102, 103
mabouya 102
multifasciata 102
quinquetaeniata 99–103
striata 99–103
Marchantia polymorpha 317
Masticophis flagellum 285
Megistolotis 261
lignarius 264
Mephitis mephitis 358
Microtus 87, 89, 91
townsendi 89
Miscanthus sinensis 305
Mixophyes fasciolatus 261, 264
Mus 284
musculus 88, 91, 284
musculus domesticus 295
musculus musculus 295
Naja nivea 170, 172
Natalobatrachus sp. 264
Natrix 331
natrix 136, 137
Neobatrachus pictus 264
Nerodia rhombifera 42–45
Nesomantis thomasseti 3
Notaden bennetti 264
melanoscaphus 258, 264
nichollsi 264
spp. 261
Nothofagus 316
Notobatrachus 350
Notophthalmus viridescens 46–48, 50, 141, 144, 154, 155
Nyctimystes sp. 1 264
sp. 2 264
Oreophryne fistulans 264
insulana 264
Osornophryne 3
buboniformis 3
percrassa 3
Palmatorappia solomonis 264
Pandanus 68
Pelobates 277
fuscus 264
varaldi 277–279
Pelodytes punctatus 264
Pelusios 370
Peromyscus 91
leucopus 118
maniculatus 91
Phrynobatrachus natalensis 264
Phrynohyas sp. 264
Phrynomantis 258
humicola 264
lateralis 264
robusta 264
stictogaster 258, 262, 264
Phrynosoma 202
Phyllobates 178, 179
aurotaenia 178, 179
chalceus 179
lugubris 178, 179
ridens 180
vittatus 178, 179
Phyllomedusa dachnicolor 277, 278
sp. 264
trinitatus 276, 278
Phyllostachys bambusoides 305
Physalaemus nattereri 137
Physignathus gilberti 284
leseuri 284
Pinus palustris 353
Pipa parva 350
pipa 277, 278
Pituophis 126, 225, 326
melanoleucus 120, 126, 225–230, 327–332
Plagioschila heterodonta 317
Platymantis papuensis 264
sp. 264
Platysaurus 312
guttatus 311–313
guttatus guttatus 311
guttatus intermedius 311
sp. 311
Plethodon 7, 10, 14, 51, 59, 113, 114, 271, 272
cinereus 14, 51, 59, 155, 271, 274
dorsalis 51, 59
glutinosus 6–15, 271–274
jordani 14, 271
nettingi 155
Pleurodema 2, 3, 316
bufonina 2
thaul 2, 317
Podocnemis expansa 158
Procyon lotor 27, 358
Pseudacris 234
brachyphona 277, 278
triseriata 234, 235, 237–242
Pseudemys 327
concinna 331
scripta 158, 327, 331–333
spp. 359
Pseudis paradoxa 264
Pseudocordylus 312
microlepidotus 311–313
Pseudonaja textilis 285
Pseudophryne bibroni 264
coriacea 264
guentheri 264
occidentalis 264
semimarmorata 264
Pseudotriton ruber 78, 85
Pteridium aquilinum 353
Ptychadena anchietae 258, 264
Pyxicephalus sp. 264
Quercus incana 353
laevis 353
margaretta 353
pumila 353
serrata 305
Ramphotyphlops 215, 216
braminus 215–220
Rana 234, 277, 279, 296, 300, 365
areolata 296–300

catesbeiana 46, 276, 278, 299
esculenta 257, 258, 260, 261
fuscigula 264
grisea 257, 258, 264
papua 264
pipiens 47, 234, 241, 242, 277–279, 296, 297, 299, 300
ridibunda 368
sphenocephala 296–300
sylvatica 234–241, 299
vitigera 264
Ranidella riparia 264
signifera 264
Regina 106
alleni 92
Rhacophorus leucomystax 264
Rheobatrachus silus 264
Rhinoderma 316
darwini 264, 317
Rhinophrynus 275–279
dorsalis 264, 275–279
Rhododendron maximum 79
Rhombophryne testudo 264
Rhyacotriton olympicus 78
Salamandra 154
Sauromalus obesus 327
Scaphiopus 277
couchi 234, 277–279
hammondi 234
holbrooki 234
sp. 264
Sceloporus 127, 129–132, 201–203, 205–211
acanthinus 203, 208
adleri 203
aeneus 102, 202, 203, 207
aeneus aeneus 206
asper 203
bulleri 204
carinatus 203
cautus 204
chrysostictus 203
clarki 203, 207, 208
couchi 204
cozumelae 204
cryptus 205
cyanogenys 204
dugesi 204
edwardtaylori 204
exsul 204
formosus 203, 211
gadoviae 203
goldmani 203
graciosus 203, 208, 310
grammicus 203
hartwegi 203
heterolepis 203
horridus 203
hunsakeri 204
insignis 204
internasalis 203
jalapae 203
jarrovi 204, 207
licki 204
lunaei 203, 208
lundelli 204
maculosus 205
madougalli 204
magister 203, 208, 310
malachiticus 203, 208
megalepidurus 204
melanorhinus 203
merriami 203, 310
mucronatus 204
nelsoni 203
occidentalis 204, 208, 242
ochoterenae 203
olivaceus 202, 203
orcutti 204, 208
ornatus 204
parvus 204
poinsetti 204
prezygus 204
pyrocephalus 203
rufidorsum 204
scalaris 102, 203, 206, 210
serrifer 204
shannonorum 203
siniferus 203
spinosus 203, 211
squamosus 203
stejnegeri 203
subpictus 204
taeniatus 203
tanneri 203
teapensis 204
torquatus 204, 211
undulatus 127–131, 204, 208
utiformis 205
variabilis 202, 204
virgatus 102, 204, 206, 208
woodi 204
Sistrurus catenatus 168
Smilisca sordida 277, 278
Solenopsis saevissima 358
Somuncuria 2, 3
somuncurensis 2
Sooglossus gardineri 3
sechellensis 3
Sorex vagrans 89
Sphenodon punctatus 60, 61, 64
Sphenomorphus quoyi 284
Sphenophryne fryi 264
robusta 264
Sternotherus 370
odoratus 33
Syrrhopus areolatus 179, 187
molinoi 175, 180, 187
Takydromus 305
takydromoides 305–310
Taudactylus diurnus 264
Telmatobufo 316
Testudo 118, 364
graeca 33, 327, 329, 332

Tetradactylus 312
seps 311–314
Thamnophis 87, 169
elegans 87, 88, 91, 92, 331
ordinoides 92
radix 87, 331
sirtalis 25, 87, 91, 92, 331
Tiliqua scincoides 60–65
Tracheloptychus 312
Trionyx ferox 359
muticus 72, 158
spp. 357
Triturus 154, 155
alpestris 47
cristatus 51
Typhlina 216
Typhlonectes compressicauda 253
Typhlops 216, 218
braminus 218, 219
lankaensis 219, 220
malcolmi 219
tenebrarum 219
veddae 219
violaceus 219
Uma notata 327
Urocyon cinereoargenteus 358
Urosaurus ornatus 103
Uta stansburiana 132, 310, 364
Varanus 266
Vermicella annulata 286
Vierraella 350
Vipera 165, 168
aspis 168
berus 168
lebetina 167
xanthina 165–169
Xenobatrachus giganteus 264
rostratus 264
Xenopus 277
laevis 145, 155, 264, 276–279
muelleri 264
tropicalis 277, 278
Xenorhina oxycephala 264
Yucca 127, 128
glauca 128–130
Zonosaurus 312

TOPICAL INDEX
HERPETOLOGICA
Volume 36 December 1980

ACTIVITY, *Elaphe obsoleta, Coluber constrictor* (relative to thermal characteristics of a cave hibernaculum) 20–26; *Desmognathus fuscus, D. ochrophaeus, Eurycea bislineata* (nocturnal surface activity) 109–115; *Gopherus agassizii* (activity correlated with season and rainfall) 301–304; *Gopherus polyphemus* (activity patterns) 353–361.

ADAPTATION, *Tiliqua scincoides* (dermal vascular system possibly functions as heat exchanger) 60–66; *Dermophis mexicanus* (life history adaptations to viviparity) 244–256; *Acanthophis antarcticus* (ambush foraging) 281–289.

AGGRESSION, *Elaphe obsoleta* (male combat) 120–127; *Colostethus trinitatis* (social behavior) 189–199.

ALBUMINS, *Rhinophrynus dorsalis* (relationships to 9 other anuran families) 275–280; *Alligator mississippiensis* (biochemical variation) 289–296.

ALLELE FREQUENCIES, *Alligator mississippiensis* (electrophoretic variation at 31 loci) 289–296.

ANATOMY, *Nerodia rhombifera* (ontogenetic changes in skull proportions) 42–46; *Eurycea* (5 spp.) (cloacal anatomy similar in non-neotenic species but differs from *E. tynerensis*) 51–60; *Tiliqua scincoides* (dermal vascular system) 60–66; *Calotes versicolor* (cytology of pineal and testes) 116–119; *Vipera xanthina* (hemipenis) 165–170; microhylid frogs (cutaneous muscles) 256–264.

BEHAVIOR, amplexus in sooglossid frogs, review of amplectic behavior in frogs 1–5; *Elaphe obsoleta, Coluber constrictor* (hibernation) 20–26; *Alligator mississippiensis* (nest attendance, agonistic behavior) 27–32; *Dendrobates pumilio* (courtship, oviposition) 69–71; *Thamnophis elegans* (prey handling) 87–93; *Desmognathus* (2 spp.), *Eurycea bislineata* (foraging) 109–115; *Elaphe obsoleta* (communication, male combat behavior) 120–127; *Bombina bombina, B. variegata* (interpopulational variation in defensive responses) 133–137; *Hemidactylium scutatum* (communal nesting, egg brooding) 141–144; *Ambystoma tigrinum* (feeding) 144–158; *Vipera xanthina* (courtship, copulation) 165–170; *Colostethus trinitatis* (social behavior, communication, parental care, color change) 189–199; *Iguana iguana* (assertion display) 221–224; *Pituophis melanoleucus* (tongue flicking) 225–231; *Plethodon glutinosus* (spatial distribution) 271–275; *Gopherus agassizii* (construction of water catchments) 301–304; *Centrolenella fleischmanni* (territoriality, reproduction) 318–326.

BIOCHEMICAL VARIATION, *Alligator mississippiensis* (electrophoresis) 289–296; *Bolitoglossa franklini, B. resplendens* (hybridization shown by electrophoresis) 335–345.

BLOOD, marine turtles (sampling methods) 17–20.

BROODING, *Hemidactylium scutatum* (egg brooding increased survival) 141–144; *Colostethus trinitatis* (egg brooding) 196–197.

CELLS, *Calotes versicolor* (cytological cycles of pineal, testes) 116–119.

CHEMOSENSATION, *Pituophis melanoleucus* (tongue flicking) 225–231.

CHROMOSOMES, *Rana areolata, R. sphenocephala, R. pipiens* (karyotypes) 296–300; cordylid evolution (chromosome evolution) 311–316; *Ascaphus truei* (karyotype) 346–352.

COLORATION, *Mabuya striata, M. quinquetaeniata* 99–104; *Colostethus trinitatis* (communication by color change) 189–199; *Bothrops schlegeli* (neonates) 231–233.

COMMUNAL, *Hemidactylium scutatum* (communal nesting) 141–144.

CONVERGENCE, *Acanthophis antarcticus* (ecology convergent with viperids) 281–289.

COURTSHIP, *Dendrobates pumilio* 69–71; *Vipera xanthina* 165–170; *Colostethus trinitatis* 189–199; *Bothrops schlegeli* (courtship, copulation) 231–233; *Centrolenella fleischmanni* 318–326; *Gopherus polyphemus* 353–361.

CRITICAL THERMAL MAXIMUM, 5 species of anurans (correlations of geographic distribution and CTM, breeding time and CTM, starvation and CTM) 234–244.

CYTOLOGY, *Calotes versicolor* (pineal, testes) 116–119.

DENSITY, *Plethodon glutinosus* (estimated by mark-recapture) 6–16; *Bombina variegata* (estimated by mark-recapture) 365–370.

DEVELOPMENT, *Caretta caretta* (sex differentiation) 33–36, (egg mortality) 72–77; *Gyrinophilus porphyriticus* (larval development) 78–86; *Eurycea quadridigitata* (larval development) 138–140; *Centrolenella fleischmanni* (egg development) 318–326.

DIET, *Thamnophis elegans* (small mammals) 87–93; *Desmognathus* (2 spp.), *Eurycea bislineata* (dietary overlap) 109–115; *Hemidactylium scutatum* (oophagy) 143; *Acanthophis antarcticus* (convergent with viperids) 281–289.

DIGGING, *Gopherus agassizii* (construction of water catchments) 301–304.

DIMORPHISM, *Mabuya striata, M. quinquetaeniata* (males larger than females) 99–104; *Dermophis mexicanus* (no sexual dimorphism) 244–256;

Acanthophis antarcticus (females larger than males) 281–289.
DISPLAY, *Iguana iguana* (assertion) 221–224.
DISTRIBUTION MAPS, *Bufo stejnegeri* (Korea) 37, 41; *Anolis barahonae* subspp. (Hispaniola) 93–98; *Eleutherodactylus gaigeae, E. chalceus, E. ridens, E. roseus, E. subsigillatus, E. taeniatus* (South America) 179–183; *Eretmochelys imbricata* (rookeries in northeastern Australia) 265–271.
DNA, cordylid lizards (great amount for lizards) 311–316.
DRINKING, *Gopherus agassizii* (construction of water catchments) 301–304.
ECOLOGY, *Plethodon glutinosus* (geographic variation in population ecology) 6–16; *Elaphe obsoleta, Coluber constrictor* (thermal ecology in a hibernaculum) 20–26; *Alligator mississippiensis* (nest attendance) 27–32; *Dendrobates pumilio* (behavioral ecology) 69–71; *Caretta caretta* (egg mortality) 72–77; *Gyrinophilus porphyriticus* (larval population ecology) 78–86; *Thamnophis elegans* (feeding ecology) 87–93; *Mabuya striata, M. quinquetaeniata* (reproductive ecology) 99–104; *Desmognathus* (2 spp.), *Eurycea bislineata* (feeding ecology, diet overlap) 109–115; *Sceloporus undulatus, Holbrookia maculata* (microhabitat, home range) 127–132; *Eurycea quadridigitata* (larval population ecology) 138–140; *Hemidactylium scutatum* (reproductive ecology) 141–144; *Ambystoma tigrinum* (feeding ecology) 144–158; *Chelydra serpentina* (nesting migrations) 158–162; *Chelonia mydas* (growth) 162–165; *Colostethus trinitatis* (behavioral ecology) 189–199; *Dermophis mexicanus* (reproductive ecology, population structure) 244–256; *Eretmochelys imbricata* (nesting ecology) 265–271; *Plethodon glutinosus* (spatial distribution) 271–275; *Acanthophis antarcticus* (food habits, reproduction, growth) 281–289; *Gopherus agassizii* (activity, construction of water catchments) 301–304; *Takydromus tachydromoides* (reproduction, growth) 305–310; *Centrolenella fleischmanni* (spatial, reproductive ecology) 318–326; *Gopherus polyphemus* (reproductive ecology) 353–361; *Bombina variegata* (movements, abundance) 365–370.
EGGS, *Plethodon glutinosus* (size) 6–16; *Caretta caretta* (incubated at various temperatures) 33–36; *Dendrobates pumilio* (incubation time) 70; *Caretta caretta* (rain-induced mortality) 72–77; *Hemidactylium scutatum* (communal nesting, brooding, survival, oophagy) 141–144; *Colostethus trinitatis* (brooding) 196–197; *Eretmochelys imbricata* (nesting biology) 265–271; *Centrolenella fleischmanni* (clutch size, survivorship, development) 318–326; *Gopherus polyphemus* (egg size, hatching success, incubation period) 353–361.
ELECTROPHORESIS, *Bitis arietans, Naja nivea* (variation in results due to storage of venom) 170–174; *Alligator mississippiensis* (biochemical variation) 289–296; *Bolitoglossa franklini, B. resplendens* (hybridization) 335–345.
EVOLUTION, *Hemidactylium scutatum* (communal nesting, solitary brooding) 141–144; *Sceloporus* (viviparity) 201–215; microhylid frogs (variation in cutaneous muscles, relation to ranoids) 256–264; *Rhinophrynus dorsalis* (immunological relationships) 275–280; cordylid lizards (chromosome evolution) 311–316; *Bolitoglossa franklini, B. resplendens* (hybridization and its evolutionary implications) 335–345.
FORAGING, *Desmognathus* (2 spp.), *Eurycea bislineata* (synchronous foraging, dietary overlap) 109–115; *Ambystoma tigrinum* (sit-and-wait strategy) 156–157; *Pituophis melanoleucus* (tongue flicking during searching) 225–231.
FUNCTIONAL MORPHOLOGY, *Tiliqua scincoides* (heat exchange in dermal vascular system) 60–66; microhylid frogs (cutaneous muscles) 256–264.
GENETICS, *Alligator mississippiensis* (low level of genetic variation) 289–296; *Bolitoglossa franklini, B. resplendens* (hybridization to early backcross generations) 335–345.
GEOGRAPHIC LOCALITIES
Australia, *Chelonia mydas* 162–165; *Eretmochelys imbricata* 265–271; *Acanthophis antarcticus* 281–289.
Bulgaria, *Bombina bombina* 133–137; *Bombina variegata* 133–137, 265–370.
California, *Gopherus agassizii* 301–304.
Chile, *Insuetophrynus acarpicus* 316–318.
Columbia, *Eleutherodactylus* spp. 175–189.
Costa Rica, *Dendrobates pumilio* 69–71; *Iguana iguana* 221–224.
Czechoslovakia, *Bombina bombina, B. variegata* 133–137.
Florida, *Alligator mississippiensis* 27–32, 289–296.
Georgia, *Caretta caretta* 33–36, 72–77; *Gopherus polyphemus* 353–361.
Guatemala, *Dermophis mexicanus* 244–256; *Bolitoglossa* spp. 335–345.
Hispaniola, *Anolis barahonae* 93–98.
Honduras, *Bothrops schlegeli* 231–233.
India, *Calotes versicolor* 116–119.
Indiana, *Eurycea bislineata* 51–60.
Iowa, *Ambystoma tigrinum* 144–158.
Japan, *Takydromus tachydromoides* 305–310.
Korea, *Bufo stejnegeri* 37–41.
Louisiana, *Nerodia rhombifera* 42–46; *Eurycea quadridigitata* 51–60; *Elaphe obsoleta* 120–127; *Alligator mississippiensis* 289–296.
Maryland, *Plethodon glutinosus* 6–16.
Mexico, *Bolitoglossa* spp. 335–345.
Missouri, *Elaphe obsoleta, Coluber constrictor* 20–26.
Nebraska, *Sceloporus undulatus, Holbrookia maculata* 127–132.
Nevada, *Gopherus agassizii* 301–304.
North Carolina, *Eurycea bislineata, E. junaluska* 51–60; *Gyrinophilus porphyriticus* 78–

86; *Rana sylvatica* 234–244; *Plethodon glutinosus* 271–275; *Rana areolata, R. sphenocephala* 296–300.
Ohio, *Desmognathus fuscus, D. ochrophaeus, Eurycea bislineata* 109–115; *Elaphe obsoleta* 120–127.
Oklahoma, *Eurycea multiplicata, E. tynerensis* 51–60; *Elaphe obsoleta* 120–127.
Ontario, *Chelydra serpentina* 158–162.
Panama, *Centrolenella fleischmanni* 318–326.
Pennsylvania, *Plethodon glutinosus* 6–16.
Poland, *Bombina bombina, B. variegata* 133–137.
Seychelles Islands, sooglossid frogs 1–5; *Ramphotyphlops braminus* 215–221.
Solomon Islands, *Cyrtodactylus* ***biordinis*** 66–69.
South Carolina, *Eurycea quadridigitata* 138–140; 4 species of anurans 234–244; *Alligator mississippiensis* 289–296.
Tennessee, *Eurycea bislineata, E. junaluska* 51–60.
Trinidad, *Colostethus trinitatis* 189–199.
Vancouver Island, *Thamnophis elegans* 87–93.
Virginia, *Hemidactylium scutatum* 141–144.
West Virginia, *Notophthalmus viridescens* 46–50.
Zambia, *Mabuya striata, M. quinquetaeniata* 99–104.

GLANDS, *Eurycea* (5 spp.) (cloacal glands) 51–60; *Calotes versicolor* (relationship of pineal and testicular cycles) 116–119.

GROWTH, *Plethodon glutinosus* (determined by mark-recapture) 6–16; *Gyrinophilus porphyriticus* (determined by size-frequency distributions of larvae) 78–86; *Eurycea quadridigitata* (larval and post-larval growth determined by size-frequency distributions) 138–140; *Chelonia mydas* (determined by mark-recapture under natural conditions) 162–165; *Dermophis mexicanus* (fetal and juvenile growth determined by size-frequency distributions) 244–256; *Acanthophis antarcticus* (determined by size-frequency distributions) 281–289; *Takydromus tachydromoides* (determined by mark-recapture, relation to reproduction) 305–310.

HABITAT, *Sceloporus undulatus, Holbrookia maculata* (microhabitat differences) 127–132; *Sceloporus* (correlates of viviparity) 201–215; *Ramphotyphlops braminus* (Seychelles Islands) 215–221.

HEART RATE, 7 species of terrestrial snakes 326–335.

HOME RANGE, *Sceloporus undulatus, Holbrookia maculata* (size, spatial distribution in both sexes) 127–132.

HYBRIDIZATION, *Bolitoglossa franklini, B. resplendens* (electrophoresis) 335–345.

IMMUNOLOGY, *Rhinophrynus dorsalis* (relationships to 9 other anuran families) 275–280.

KARYOTYPE, *Rana areolata, R. sphenocephala* (comparisons with *R. pipiens*) 296–300; cordylid lizards (chromosome evolution) 311–316; *Ascaphus truei* (systematic relationship) 346–352.

KEY, *Cyrtodactylus* (key to species from Solomon Islands) 68–69.

LARVAE, *Gyrinophilus porphyriticus* (variable growth rates and larval period) 78–86; *Eurycea quadridigitata* (growth and larval period) 138–140; 5 species of anurans (thermal tolerance of tadpoles) 234–244; *Insuetophrynus acarpicus* (description of tadpole) 316–318.

LIFE HISTORY, *Plethodon glutinosus* (geographic variation in population parameters) 6–16; *Gyrinophilus porphyriticus* (variable larval period) 78–86; *Mabuya striata, M. quinquetaeniata* (reproduction, size at sexual maturity) 99–104; *Eurycea quadridigitata* (larval period, size at sexual maturity) 138–140; *Chelonia mydas* (immature growth and age at maturity) 162–165; *Dermophis mexicanus* (age at maturity, growth, reproductive biology, population structure, sex ratio) 244–256; *Takydromus tachydromoides* (growth rate, size at maturity) 305–310.

MARK-RECAPTURE, *Plethodon glutinosus* (population study) 6–16; *Chelydra serpentina* (nesting migrations) 158–162; *Chelonia mydas* (growth) 162–165; *Colostethus trinitatis* (social behavior) 189–199; *Centrolenella fleischmanni* (movements, reproductive behavior) 318–326; *Bombina variegata* (movement ecology) 365–370.

METABOLISM, *Caiman crocodilus* (standard metabolic rate, specific dynamic action, relation to nutritional status) 361–364.

METAMORPHOSIS, 5 species of anurans (thermal tolerance) 234–244.

METHODS, marine turtles (sampling blood, cerebrospinal fluid) 17–20; *Bitis arietans, Naja nivea* (variation in electrophoretic properties due to venom storage) 170–174; *Rana areolata, R. sphenocephala* (obtaining karyotypes from tadpoles) 296–300.

MORPHOLOGY, *Bolitoglossa franklini, B. resplendens* (expression of hybridization) 335–345.

MOVEMENT, *Chelydra serpentina* (nesting migrations) 158–162; *Bombina variegata* (movement ecology, abundance) 365–370.

MUSCULATURE, microhylid frogs (phylogenetic and functional significance of cutaneous muscles) 256–264.

NEST ATTENDANCE, *Alligator mississippiensis* (agonistic behavior) 27–32.

NESTING, *Eretmochelys imbricata* (density, clutch size, incubation success, nest destruction) 265–271; *Gopherus polyphemus* (timing, location, nest predation) 353–361.

ONTOGENY, *Nerodia rhombifera* (skull proportions) 42–46; *Mabuya striata, M. quinquetaeniata* (color patterns) 99–104.

PARTHENOGENESIS, *Ramphotyphlops braminus* 215–221.

PHYLOGENY, sooglossid frogs (implications of amplectic behavior) 1–5; *Sceloporus* (correlates of viviparity) 201–215; microhylid frogs (significance of cutaneous muscles) 256–264; *Rhino-*

phrynus dorsalis (biochemical relationships) 275–280; cordylid lizards (chromosomal evolution and DNA) 311–316.

PHYSIOLOGY, *Tiliqua scincoides* (blood flow in dermal vascular system during heating and cooling) 60–66; *Iguana iguana* (partial control of assertion display) 221–224; 7 species of terrestrial snakes (heart rate-ventilatory response) 326–335; *Caiman crocodilus* (metabolic rate in recently fed and fasted animals) 361–364.

PLACENTA, *Sceloporus* (evolution of viviparity) 201–215.

POPULATION STRUCTURE, *Plethodon glutinosus* (geographic variation) 6–16; *Dermophis mexicanus* (size-frequency distributions) 244–256; *Eretmochelys imbricata* (size of nesting females) 265–271; *Bombina variegata* (sex ratio) 365–370.

PREDATION, *Thamnophis elegans* (small mammal predation) 87–93; *Desmognathus* (2 spp.), *Eurycea bislineata* (synchronous foraging, dietary overlap) 109–115; *Bombina bombina, B. variegata* (behavioral responses to predators) 133–137; *Eretmochelys imbricata* (predation on nests) 265–271; *Gopherus polyphemus* (predation on nests) 353–361.

RADIOTELEMETRY, *Chelydra serpentina* (nesting migrations) 158–162.

REPRODUCTION, *Plethodon glutinosus* (clutch size, egg diameter) 6–16; *Dendrobates pumilio* (courtship, oviposition, clutch size) 69–71; *Mabuya striata, M. quinquetaeniata* (times, modes, coincidence with climatic factors, fat bodies, clutch sizes) 99–104; *Calotes versicolor* (relationship of pineal and testicular cycles) 116–119; *Sceloporus* (evolution of viviparity) 201–215; *Ramphotyphlops braminus* (parthenogenesis) 215–221; *Bothrops schlegeli* (captive pair) 231–233; *Dermophis mexicanus* (cycling, clutch size, fertilization, gestation, birth) 244–256; *Eretmochelys imbricata* (nesting biology) 265–271; *Acanthophis antarcticus* (cycling, litter sizes) 281–289; *Takydromus tachydromoides* (relation to growth) 305–310; *Centrolenella fleischmanni* (territory, mating success) 318–326; *Gopherus polyphemus* (courtship, nesting, clutch size, low reproductive rate) 353–361.

SEX DIFFERENTIATION, *Caretta caretta* (eggs incubated at various temperatures) 33–36.

SIZE, *Gyrinophilus porphyriticus* (larval period model based on size-frequency distributions) 78–86; *Thamnophis elegans* (size of prey) 87–93.

SPATIAL DISTRIBUTION, *Sceloporus undulatus, Holbrookia maculata* (home ranges) 127–132; *Plethodon glutinosus* (individual interaction) 271–275; *Centrolenella fleischmanni* (territorial spacing) 318–326; *Bombina variegata* (movement ecology) 365–370.

SURVIVORSHIP, *Plethodon glutinosus* (determined from age and sex distributions) 6–16; *Caretta caretta* (rain-induced mortality of eggs) 72–77; *Hemidactylium scutatum* (egg brooding increased embryonic survival) 141–144; *Centrolenella fleischmanni* (eggs) 318–326.

SYNONYMY, *Bufo kangi* (synonym of *B. stejnegeri*) 37–41; *Eleutherodactylus areolatus* (synonym of *E. chalceus*), *Syrrhopus molinoi* (synonym of *E. ridens*) 175–189.

SYSTEMATICS, sooglossid frogs (amplectic behavior, phylogeny) 1–5; *Bufo stejnegeri* (redescription and relationship to *B. bufo* group) 37–41; *Nerodia rhombifera* (effects of ontogeny on skull proportions, use in taxonomy) 42–46; *Cyrtodactylus* ***biordinis*** (sp. nov.) 66–69; *Anolis barahonae* (subspp. nov. with comparisons to other subspp.) 93–98; *Eleutherodactylus* spp. (redescriptions, synonomies, distribution) 175–189; *Ramphotyphlops braminus* (distribution, variation) 215–221; microhylid frogs (phylogenetic significance of cutaneous muscles) 256–264; *Rhinophrynus dorsalis* (biochemical relationship to 9 other anuran families) 275–280; *Alligator mississippiensis* (biochemical variation) 289–296; *Rana areolata, R. sphenocephala* (cytotaxonomy) 296–300; cordylid lizards (chromosomal evolution, DNA) 311–316; *Bolitoglossa franklini, B. resplendens* (hybridization) 335–345; *Ascaphus truei* (relationship based on karyotype) 346–352.

TAXONOMY, *Bufo stejnegeri* (redescription, relationship to *B. bufo* group) 37–41; *Nerodia rhombifera* (effects of ontogeny on skull proportions) 42–46; *Cyrtodactylus* ***biordinis*** (sp. nov.) 66–69; *Anolis barahonae* (subspp. nov. with comparisons to other subspp.) 93–98; *Eleutherodactylus* spp. (redescriptions, synonomies, distribution) 175–189; *Ramphotyphlops braminus* (nomenclature) 215–221; *Rana* (cytotaxonomy of 3 species) 296–300; *Ascaphus truei* (karyology, relationship to other anurans) 346–352.

TERRITORY, *Colostethus trinitatis* (size, defense) 189–199; *Centrolenella fleischmanni* (relation to mating success) 318–326.

THERMAL ACCLIMATION, 5 species of anurans (during development and metamorphosis) 234–244.

THERMAL GRADIENT, *Elaphe obsoleta, Coluber constrictor* (in a cave hibernaculum) 20–26.

THERMOREGULATION, *Tiliqua scincoides* (functional morphology of dermal vascular system) 60–66.

TONGUE, *Pituophis melanoleucus* (rate of tongue flicking) 225–231.

TUMOR, *Notophthalmus viridescens* (neuroblastoma) 46–50.

VENOM, *Bitis arietans, Naja nivea* (change in electrophoretic properties due to storage) 170–174.

VENTILATION, 7 species of terrestrial snakes (similar to aquatic snakes) 326–335.

VIVIPARITY, *Sceloporus* (environmental, geographic, phylogenetic, life history correlates) 201–215; *Dermophis mexicanus* (reproductive biology) 244–256; *Acanthophis antarcticus* (reproductive biology) 281–289.

VOCALIZATION, *Colostethus trinitatis* 189–199; *Centrolenella fleischmanni* 318–326.

HERPETOLOGICA

Volume 36 December 1980

TABLE OF CONTENTS

Adams, Susan E., Michael H. Smith, and Ramone Baccus
Biochemical Variation in the American Alligator 289
Announcement: Passing of Arthur Loveridge 200
Annual Financial Statement 107
Antonio, Frederick B.
Mating Behavior and Reproduction of the Eyelash Viper (*Bothrops schlegeli*) in Captivity 231
Baccus, Ramone (see Adams, Susan E.)
Bachmann, Marilyn D. (see Lindquist, Sarah B.)
Bajger, Jan
Diversity of Defensive Responses in Populations of Fire Toads (*Bombina bombina* and *Bombina variegata*) 133
Barker, David G. (see Murphy, James B.)
Bell, Rebecca (see Kraemer, Jed E.)
Beshkov, Vladimir A., and David L. Jameson
Movement and Abundance of the Yellow-Bellied Toad *Bombina variegata* 365
Bogart, James P. (see Green, David M.)
Brooks, Ronald J. (see Obbard, Martyn E.)
Brown, Walter C., and Michael McCoy
A New Species of Gecko of the Genus *Cyrtodactylus* from Guadalcanal Island, Solomon Islands 66
Bruce, Richard C.
A Model of the Larval Period of the Spring Salamander, *Gyrinophilus porphyriticus*, Based on Size-Frequency Distributions 78
Burton, Thomas C.
Phylogenetic and Functional Significance of Cutaneous Muscles in Microhylid Frogs 256
Bury, R. Bruce (see Medica, Philip A.)
Chiszar, David, Valerie Lipetz, Kent Scudder, and Elizabeth Passanello
Rate of Tongue Flicking by Bull Snakes and Pine Snakes (*Pituophis melanoleucus*) During Exposure to Food and Non-Food Odors 225
Corcoran, Michael F., and Joseph Travis
A Comparison of the Karyotypes of the Frogs *Rana areolata, Rana sphenocephala*, and *Rana pipiens* 296
Counts, Clement L., III
A Neural Neoplasm in the Eastern Newt, *Notophthalmus viridescens* 46
Cullom, Shelby J., and Albert Schwartz
Variation in the Hispaniolan Giant Anole *Anolis barahonae* with the Description of Two New Subspecies 93
Cupp, Paul V., Jr.
Thermal Tolerance of Five Salientian Amphibians During Development and Metamorphosis 234
Daugherty, Charles H. (see Green, David M.)
Daugherty, Charles H. (see Maxson, Linda R.)
Diaz, N. F. (see Formas, J. R.)
Drane, C. R., and C. J. W. Webb
Functional Morphology of the Dermal Vascular System of the Australian Lizard *Tiliqua scincoides* 60
Droge, Dale L. (see Jones, Steven M.)
Fitzgerald, Kevin T. (see Guillette, Louis J.)
Formas, J. R., N. F. Diaz, and J. Valencia
The Tadpole of the Chilean Frog *Insuetophrynus acarpicus* 316
Garner, James A. (see Landers, J. Larry)

Gatten, Robert E., Jr.
Metabolic Rates of Fasting and Recently Fed Spectacled Caimans (*Caiman crocodilus*) 361
Gill, Douglas E. (see Harris, Reid N.)
Gillingham, James C.
Communication and Combat Behavior of the Black Rat Snake (*Elaphe obsoleta*) 120
Green, David M., Charles H. Daugherty, and James P. Bogart
Karyology and Systematic Relationships of the Tailed Frog *Ascaphus truei* 346
Greer, Beverly J., and Kentwood D. Wells
Territorial and Reproductive Behavior of the Tropical American Frog *Centrolenella fleischmanni* 318
Gregory, Patrick T., Malcolm Macartney, and Donald H. Rivard
Small Mammal Predation and Prey Handling Behavior by the Garter Snake *Thamnophis elegans* 87
Guillette, Louis J., Jr., Richard E. Jones, Kevin T. Fitzgerald, and Hobart M. Smith
Evolution of Viviparity in the Lizard Genus *Sceloporus* 201
Haldar, C. (see Misra, C. nee Haldar)
Harris, Reid N., and Douglas E. Gill
Communal Nesting, Brooding Behavior, and Embryonic Survival of the Four Toed Salamander *Hemidactylium scutatum* 141
Hattingh, J. (see Willemse, G. T.)
Hazlett, Brian A.
Temporal Pattern of the Assertion Display of the Lizard *Iguana iguana* 221
Herpetologists' League Student Award 106
Holomuzki, Joseph R.
Synchronous Foraging and Dietary Overlap of Three Species of Plethodontid Salamanders 109
Hunt, Sue R. (see Sexton, Owen J.)
Index to Scientific Names for Volume 36 372
Jacob, James S.
Heart Rate-Ventilatory Response of Seven Terrestrial Species of North American Snakes 326
Jameson, David L. (see Beshkov, Vladimir A.)
Jones, Richard E. (see Guillette, Louis J.)
Jones, Steven M., and Dale L. Droge
Home Range Size and Spatial Distributions of Two Sympatric Lizard Species (*Sceloporus undulatus, Holbrookia maculata*) in the Sand Hills of Nebraska 127
Kraemer, Jed E., and Rebecca Bell
Rain-Induced Mortality of Eggs and Hatchlings of Loggerhead Sea Turtles (*Caretta caretta*) on the Georgia Coast 72
Kushlan, James A., and Marilyn S. Kushlan
Function of Nest Attendance in the American Alligator 27
Kushlan, Marilyn S. (see Kushlan, James A.)
Landers, J. Larry, James A. Garner, and W. Alan McRae
Reproduction of Gopher Tortoises (*Gopherus polyphemus*) in Southwestern Georgia 353
Limerick, Sandra
Courtship Behavior and Oviposition of the Poison-Arrow Frog *Dendrobates pumilio* 69
Limpus, Colin J.
Observations on the Hawksbill Turtle (*Eretmochelys imbricata*) Nesting Along the Great Barrier Reef 265
Limpus, Colin J., and David G. Walter
The Growth of Immature Green Turtles (*Chelonia mydas*) Under Natural Conditions 162
Lindquist, Sarah B., and Marilyn D. Bachmann
Feeding Behavior of the Tiger Salamander, *Ambystoma tigrinum* 144
Lipetz, Valerie (see Chiszar, David)
Luckenbach, Roger A. (see Medica, Philip A.)
Lynch, John D.
Systematic Status and Distribution of Some Poorly Known Frogs of the Genus *Eleutherodactylus* from the Chocoan Lowlands of South America 175
Macartney, Malcolm (see Gregory, Patrick T.)

Matsui, Masafumi
The Status and Relationships of the Korean Toad, *Bufo stejnegeri* Schmidt 37
Maxson, Linda R., and Charles H. Daugherty
Evolutionary Relationships of the Monotypic Toad Family Rhinophrynidae: a Biochemical Perspective 275
McCoy, Michael (see Brown, Walter C.)
McRae, W. Alan (see Landers, J. Larry)
Medica, Philip A., R. Bruce Bury, and Roger A. Luckenbach
Drinking and Construction of Water Catchments by the Desert Tortoise, *Gopherus agassizii*, in the Mojave Desert 301
Misra, C. (nee Haldar), and J. P. Thapliyal
Inverse Seasonal Cytological Cycles in Testes and Pineal Organs of the Bloodsucker *Calotes versicolor* 116
Mrosovsky, N. (see Yntema, C. L.)
Murphy, James B., and David G. Barker
Courtship and Copulation of the Ottoman Viper (*Vipera xanthina*) with Special Reference to Use of the Hemipenes 165
Nomenclatural Notices 200, 280, 370
Nussbaum, Ronald A.
Phylogenetic Implications of Amplectic Behavior in Sooglossid Frogs 1
The Brahminy Blind Snake (*Ramphotyphlops braminus*) in the Seychelles Archipelago: Distribution, Variation, and Further Evidence of Parthenogenesis 215
Obbard, Martyn E., and Ronald J. Brooks
Nesting Migrations of the Snapping Turtle (*Chelydra serpentina*) 158
Odierna, Gaetano (see Olmo, Ettore)
Olmo, Ettore, and Gaetano Odierna
Chromosomal Evolution and DNA of Cordylid Lizards 311
Owens, David Wm., and Georgita J. Ruiz
New Methods of Obtaining Blood and Cerebrospinal Fluid from Marine Turtles 17
Papenfuss, Theodore J. (see Wake, David B.)
Pasanello, Elizabeth (see Chiszar, David)
Rivard, Donald H. (see Gregory, Patrick T.)
Rossman, Charles E.
Ontogenetic Changes in Skull Proportions of the Diamondback Water Snake, *Nerodia rhombifera* 42
Ruiz, Georgita J. (see Owens, David Wm.)
Schwartz, Albert (see Cullom, Shelby J.)
Scudder, Kent (see Chiszar, David)
Semlitsch, Raymond D.
Geographic and Local Variation in Population Parameters of the Slimy Salamander *Plethodon glutinosus* 6
Growth and Metamorphosis of Larval Dwarf Salamanders (*Eurycea quadridigitata*) 138
Sever, David M.
Cloacal Anatomy of Male Brook Salamanders (*Eurycea*) 51
Sexton, Owen J., and Sue R. Hunt
Temperature Relationships and Movements of Snakes (*Elaphe obsoleta, Coluber constrictor*) in a Cave Hibernaculum 20
Shine, Richard
Ecology of the Australian Death Adder *Acanthophis antarcticus* (Elapidae): Evidence for Convergence with the Viperidae 281
Simbotwe, Malumo P.
Reproductive Biology of the Skinks *Mabuya striata* and *Mabuya quinquetaeniata* in Zambia 99
Smith, Hobart M. (see Guillette, Louis J., Jr.)
Smith, Michael H. (see Adams, Susan E.)
Summary of Business Conducted at the Annual Meeting, 1979 105
Statement of Ownership, Management, and Circulation 371
Takenaka, Sen
Growth of the Japanese Grass Lizard *Takydromus tachydromoides* in Relation to Reproduction 305
Thapliyal, J. P. (see Misra, C. nee Haldar)

Topical Index for Volume 36 ---- 378
Travis, Joseph (see Corcorcan, Michael F.)
Valencia, J. (see Formas, J. R.)
Wake, David B., Suh Y. Yang, and Theodore J. Papenfuss
Natural Hybridization and Its Evolutionary Implications in Guatemalan Plethodontid Salamanders of the Genus *Bolitoglossa* ---- 335
Wake, Marvalee H.
Reproduction, Growth, and Population Structure of the Central American Caecilian *Dermophis mexicanus* ---- 244
Walter, David G. (see Limpus, Colin J.)
Webb, G. J. W. (see Drane, C. R.)
Wells, Kentwood D.
Social Behavior and Communication of a Dendrobatid Frog (*Colostethus trinitatis*) ... 189
Spatial Associations Among Individuals in a Population of Slimy Salamanders (*Plethodon glutinosus*) ---- 271
Wells, Kentwood D. (see Greer, Beverly J.)
Willemse, G. T., and J. Hattingh
Effect of Drying and Storage on Electrophoretic Properties of Venom from Puff Adders (*Bitis arietans*) and Cape Cobras (*Naja nivea*) ---- 170
Yang, Suh Y. (see Wake, David B.)
Yntema, C. L., and N. Mrosovsky
Sexual Differentiation in Hatchling Loggerheads (*Caretta caretta*) Incubated at Different Controlled Temperatures ---- 33